

The Effect of Chicken Manure and Biochar Application on The Growth and Yield of Cauliflower (*Brassica oleraceae* var. *Botritys* L)

Agostinho da C. Moniz^a, Donata O. de Araujo^b, Marcal Gusmão^c, Lucio M. Gomes^d, Natalino de Araujo^{e*}, Luis M. Branco^f, Jose A. Marcal^g

*a,b,c,d,e,f,g*Department of Agronomy, Faculty of Agriculture, Universidade Nacional Timor Lorosa'e – Untl, Dili - Timor Leste

^aEmail: natalino.dearaujo@untl.edu.tl

Abstract

Field research was carried out in Raifun village-Maliana Timor Leste from September 2023 to January 2024 at 226 meters above sea level, aiming to determine the effect of chicken manure and biochar on the growth and yield of cauliflower plants. This study used a Randomized Complete Block design (RCBD) that was repeated three times in 3 blocks. the first factor in this study is chicken manure which consists of 3 levels of treatment, namely: without chicken manure (M0), chicken manure dose of 10 t h⁻¹ (M1), and chicken manure with a dose of 20 t h⁻¹. The second factor is biochar which consists of 3 levels of treatment, including: without biochar (B0), biochar 5 t h⁻¹, and biochar 10 t h⁻¹. The results showed that there was an interaction between the two factors in the parameters of plant height, flower weight per plant, and flower weight per hectare. the highest production per hectare was obtained by a combination of a chicken manure treatment dose of 20 t h⁻¹ and a biochar dose of 10 t h⁻¹ of 19.1667 tons. The lowest production per hectare was achieved by a combination of treatments without the application of chicken manure and biochar amounting to 9.1733 tons.

Keywords: Chicken Manure; Biochar; Growth; Yield; Cauliflower.

Received: 6/12/2024

Accepted: 8/12/2024

Published: 8/22/2024

* Corresponding author.

1. Introduction

Cauliflower (*Brassica oleraceae* var. *Botritys* L.) is one of the horticultural plants categorized as vegetable plants that have white flowers. Cauliflower belongs to the family Brassicaceae [26]. In addition, cauliflower is also famous as a plant that has a high economic value compared to other vegetable crops [12]. Therefore the majority of people in the world grow and consume cauliflower as a vegetable that is rich in nutrients, vitamins, and minerals. Cauliflower thrives in countries of subtropical such as Europe, China, Japan, and Korea. The high economic value led to the spread of this plant through trade and distribution activities to African and Asian countries [12]. Cauliflower plants contain 93% water in all plant organs such as stems, leaves, and flowers. In addition, cauliflower also contains some nutrients calcium 2%, potassium 2%, phosphorus 6%, magnesium 4%, iron 3%, and fiber 2.0%, and is a source of vitamins A, B, and C. This plant is also important for human health including anticipating attacks of bone disease, heart, cancer, eye, stomach, and asthma, therefore it is important to cultivate [8]. Cauliflower plants have been cultivated in small quantities in East Timor, although they haven't yet achieved optimum production. Cauliflower cultivated in several regions of Timor Leste has an average production per hectare of only 11.95 t h⁻¹ [7]. The amount of production mentioned is still very low when compared to the average cauliflower production in other countries such as Indonesia, which is 12.08 t h⁻¹ [9]. Organic matter is a soil improvement material derived from plant and animal residues. Some organic materials that are very important for improving the physical, chemical, and biological properties of the soil are manure and biochar, chicken manure that has sufficient nutrient content is one of the manures for plant growth and yield. In addition to manure, organic matter that is rich in carbon is biochar which has a role in neutralizing soil acidity and be able to bind water in the soil, Cauliflower production that did not increase from year to year in Timor Leste is due to the absence of knowledge about the use of appropriate technology, including the use of chicken manure and biochar organic material so that production could be increased. Therefore, it is necessary to study the application of chicken manure and biochar material as one of the technologies in increasing cauliflower production.

Objectives

The objectives of this research are :

- To know the effect of chicken manure on the growth and yield of cauliflower
- To know the effect of biochar on the growth and yield of cauliflower
- To know the effect of interactions of chicken manure and biochar on the growth and yield of cauliflower.

2. Research Methods

2.1. Site and Time

An experimental trial was conducted in Maliana Sub District, Bobonaro Municipality from September

2023 to January 2024.

2.2. Material and Equipment

Materials and Equipment used in the research are the seed of cauliflower, chicken manure, biochar, line meters, analytic balance, pH tester, rules, manual balance, sword, hoe, crowbar, hammer, bucket pipe, oven, thermometer, triplex, nail, dendrometer, caliper, string of raffia.

2.3. Experimental Design

The agronomic trial was a Randomized Complete Block Design (RCBD) with two factors. The first factor was chicken manure (M), consisting of 3 treatment levels, M0= without chicken manure (0 t/ha), M1= application of chicken manure doze 10 t h⁻¹, M2= application of chicken manure doze 20 t h⁻¹. The second factor was biochar (B), consisting of 3 treatment levels, B0= Without biochar (0 t/ha), B1= application of biochar doze 5 t h⁻¹, B2= application of biochar doze 10 t h⁻¹. The treatment combination was established at 3 blocks. Each block had 9 plots. The Parameters observed were water content, the plant height, the number of leaves, the weight of flowers per plant, and the weight of flowers per hectare. Other than these above treatments, all other crop-specific were employed across treatments i.e., land preparation, crop care, and management were similar to common farming practices of Maliana farmers.

2.4. Data Analysis

The data observed had been analyzed variation (ANOVA) to indicate the interaction between 2 factors and significant differences. Furthermore, to compare each other treatments using LSD 5 % level. The media utilized in statistical analysis were software STAR 2.1 IRRI.

3. Result and Discussion

The results of ANOVA showed the interaction between chicken manure and biochar on the number of leaves at observation 2 weeks after planting, plant height, flower weight per plant, and flower weight per hectare. The result of the statistical (picture 1) indicated that applying chicken manure did not affect the soil water content, however the higher percentage of soil water content achieved on applying chicken manure 20 t h⁻¹ (42.74 %), and the lower value of soil water content at without applying chicken manure (42.44 %). Applying chicken manure 20 t h⁻¹ showed 0.11 % soil water content higher than applying chicken manure 10 t h⁻¹, and 0.30 % when compared to without applying any organic fertilizer. The higher percentage of soil water content in a higher dose of chicken manure due to the high volume of organic fertilizer can improve the structure, texture, and porosity of the soil and subsequently, the balance between water and oxygen in the soil. Appropriate soil structure and texture can bind and provide water in the soil pore thereby reducing the temperature in the soil. Evaporation is one indication of the amount of water in the soil.




Figure1: Effect of chicken manure and Biochar on soil water Content (%)

The appropriate quantity of chicken manure provides an opportunity for the microorganisms in the soil to establish the decomposition process faster, thus good physical, chemical, and biological soil conditions will improve the ability of the soil to bind water in the soil. The process of decomposition by microorganisms requires an appropriate temperature to be effectively decomposed [24]. Explained that water and temperature are primary factors in the process of decomposition of organic materials such as plant and animal residues which are directly related to microbial activity in the soil.

Picture 1 it shall be seen that applying of biochar affects the level of soil moisture, where the application of biochar of 10 t h⁻¹ has a different effect from without the application of biochar, but there is no detected significant difference with the application of biochar of 5 t h⁻¹. This is due to biochar being a soil improvement material where the process involves oxygen and converts biomass into a stable form of carbon that can be used to improve soil health and soil productivity [11]. Reveal that biochar is high solid carbon produced from the pyrolysis of biomass such as wood, manure, and leaves under high temperatures and low oxygen conditions which is used for agricultural applications as a soil amendment.

The results in Table 1 show the effect of applying chicken manure on the number of leaves per plant at 4 and 6 weeks after planting. The application of chicken manure 20 t h⁻¹ had a different effect from the without the application of chicken manure, but there was no noticeable difference with the application of chicken manure 10 t h⁻¹ at the observation of 6 weeks after planting. Observation, 4 weeks after planting showed differences in influence between each treatment. The highest number of leaves in the second week of observation showed interaction (M2B0= 9.5833), week 4 (14.75), and week 6 (31.00). Conversely, the number of leaves was less in the second week amounting to 7.1667 in the combination of M2B2, week 4 (12.17), and week 6 (23.61). The

application of the appropriate dose of chicken manure has a role in the growth of the vegetative part of the plant. These results are due to the optimal application of manure to create better physical, chemical, and biological soil conditions and subsequently absorption by roots, photosynthesis processes, and translocation to the vegetative parts of plants including the increase in the number of leaves [15]. Explained that optimal nutrition in the soil has a role in stimulating the growth of vegetative and reproductive parts of plants to provide maximum results. [4]. confirmed that the presence of organic fertilizers can affect the vegetative growth of plants, thus the final yield of plants could be achieved maximally [16].

Table 1: Effect of chicken manure and biochar on the number of leaves (sheet)

Treatments	week 2	Treatments	week 4	Growth Rates (GR)	week 6
M0B0	7.1667c	M0	12.17c	11.44	23.61b
M0B1	7.25c	M1	14.25b	16.22	30.47a
M0B2	7.333c	M2	14.75a	16.25	31a
M1B0	7.75bc	LSD	0.4856		0.9137
M1B1	8.75ab	CV	3.54		3.22
M1B2	8.25b	F pr	0		0
M2B0	9.5833a	B0	13.64A	14.89	28.53A
M2B1	8.3333b	B1	13.67A	14.19	27.86A
M2B2	9ab	B2	13.86A	14.83	28.69A
LSD:	0.8634	F pr	0.5826		0.1557
CV:	6.12				
F pr:	0.0219				

Justification: The letter followed by the same lowercase or uppercase does not show different effects at the LSD 5 %

Observations 2 weeks after planting showed that the combination of applying chicken manure 20 t h⁻¹ and without biochar was different in effect from applying chicken manure combined with biochar 10 t h⁻¹, 5 t h⁻¹, or without biochar, but there was no difference in effect with the combination of applying chicken manure 20 tons per hectare and biochar 10 t h⁻¹. The application of chicken manure and biochar in suitable soil environmental conditions can improve the physical, chemical, and biological conditions of the soil so that plant roots easily absorb nutrients for plants [9]. Explained that solid organic materials such as chicken residues and biochar provide a decrease in soil density and high organic C levels, which leads to better soil structure and subsequently plant roots are effortless to develop and optimize the absorption of nutrients and water in the soil.

Biochar application showed no significant effect on leaf count at all observation times. However, biochar with a dose of 10 t h⁻¹ gave the highest leaf count result of 28.69 no and the lowest 28.53 in treatment without biochar. Applying Biochar with the appropriate dose has the potential to improve soil conditions and plant productivity in the absorption of nutrients, water, and other minerals for further use for plant needs. The role of biochar in

revitalizing soil conditions is the effectiveness of fertilization so that biochar can bind nutrients in maximum conditions and release for the necessity of plant absorption (slow release), thus plants avoid nutrient poisoning (especially micronutrients). Other studies found that the biochar from rice husks could build soil and increase plant efficiency in utilizing nutrients for the development of plants [8].

The interaction between chicken manure and biochar at plant height (table 2) showed the highest yield in the second week at 20t/h and without biochar (23.1667 cm), at weeks 4 and 6 at 20 t h⁻¹ and 10 t h⁻¹ (39.5 cm, and 51.5 cm) biochar.

Table 2: Effect of Chicken Manure and Biochar on Plant Height (cm)

Treatments	weeks 2	GR 2-4	weeks 4	Growth Rates (GR)	weeks 6
M0B0	15.1667c	9.5833	24.75d	17.75	42.5d
M0B1	13e	9.25	22.25e	18	40.25e
M0B2	13.25d	9	22.25e	17.25	39.5e
M1B0	21.25b	12.75	34c	11.833	45.8333c
M1B1	22.5ab	13.25	35.75c	14.42	50.1667ab
M1B2	21b	13	34c	15.08	49.0833b
M2B0	23.1667a	14.17	37.3333b	13.92	51.25ab
M2B1	21.5833b	12.5	34.0833c	13.42	47.5bc
M2B2	22.75ab	16.75	39.5a	12	51.5a
LSD:	1.2172		1.9524		2.2321
CV:	3.64		3.58		2.78
Fpr:	0.0036		0.0002		0.0002

Justification: The letter followed by the same lowercase does not show a difference effect at the LSD 5 %

The rate of increase in plant height (2 mst-4mst) for the highest yield of chicken manure 20 t h⁻¹ and without biochar was 14.17 cm (61.16 %), while the growth rate of (4 wap– 6 wap) was 13.92 cm (37.29 %). The rate of increase in height for chicken manure 20 t h⁻¹ and biochar 10 t h⁻¹ was 16.75 cm (73.63%) and 12 cm (30.38%). These results showed that the rate of increase in plant height decreased after 4 weeks. This is due to in the vegetative growth phase, plants actively absorb nutrients for the formation and extension of vegetative organs, and in the phase of flowering, fruit, and seed formation, 80% of photosynthate results are translocated to fruits and seeds. The addition of chicken manure and biochar which is considered an organic material that can mobilize nutrients in the soil to form ion particles for effortless absorption [9]. [26] Added that the use of chicken manure and biochar simultaneously with appropriate quantities affects the speed of nutrient absorption by plants.

The interaction between chicken manure fertilizer and biochar is seen in production per plant, where the combination of chicken manure 20 t h⁻¹ and biochar 10 t h⁻¹) provides a stronger effect than other combinations with a yield of 0.4792 kg. The combination of applying 20 t h⁻¹ chicken manure and without biochar achieved

higher flower weight per plant (0.3849 kg) than 10 t h⁻¹ chicken manure combined with biochar 10 t h⁻¹ (0.3533 kg), 5 t h⁻¹ (0.3643 kg) and no biochar (0.3661 kg). However, there is no significant difference between the application of chicken manure 20 t h⁻¹ and biochar 5 t h⁻¹ (0.3724 kg). The lowest yield was obtained without chicken manure and biochar (0.2293 kg). From these results, it is suspected that applying chicken manure and biochar in higher doses could improve soil properties and provide nutrients to increase flower production [26].

Said that the use of organic materials such as chicken manure fertilizer and optimal biochar is very important to improve soil quality and character such as increasing the ability of the soil to bind water so that soil temperature could be controlled, increasing the ability of the soil to absorb nutrients and water, thus the photosynthesis process be able to run well and the resulting photosynthesis is utilized by plant parts such as flowers, fruits, and seeds. Research by [2] and [17] That the addition of chicken manure 20 t h⁻¹ provides higher cauliflower flower production per hectare compared to the use of lower doses due to the content of essential macronutrients in chicken manure is sufficient for the necessity of cauliflower [29]. Stated that biochar can optimize plant growth and yield, reducing nutrient loss due to leaching [23]. added that microbial activity in the soil increases with the addition of biochar in the soils, thereby increasing organic matter in the soil [18]. Revealed that the addition of biochar can increase the ability of soil to bind water, increase CEC, and provide essential nutrients to be absorbed by plants.




Figure2: Effect of chicken manure and biochar on weight of flower per plant (kg)




Figure3: Effect of chicken manure and biochar on weight of flower per hectare (tone)

The application of chicken manure 20 t h⁻¹ combined with biochar 10 t h⁻¹ with a yield of 19.1667-ton h⁻¹ is different from the effect of the combination of chicken manure fertilizer and other biochar on flower production per hectare (picture 3). The application of 20 t h⁻¹ chicken manure that is not accompanied by biochar gives better results (15.3967 tons) than the application of 20 t h⁻¹ chicken manure, 10 t h⁻¹ combined with 5 t h⁻¹ biochar and without biochar picture 3. The results mentioned above indicate that one of the roles of chicken manure and biochar which is organic matter is to be a proper habitat for the development of microorganisms [5]. Explained that the combination of organic materials such as chicken shrimp fertilizer and biochar which is used as a habitat for microorganisms can increase soil biological activities due to decreased competition between microorganisms in the soil. Increased activity of soil microorganisms causes an increase in the amount of nutrients in the soil which are further absorbed by plants for the needs of crop production, especially flowers, fruits, and seeds.

4. Conclusion

The result and discussion concluded that there was an interaction between chicken manure and biochar on plant height, flower weight per plant, and flower weight per hectare. The factors of chicken manure and biochar each significantly influence soil moisture content and leaves number. The highest flower production per hectare was obtained from a combination of chicken manure treatment doses of 20 t h⁻¹ and biochar of 10 t h⁻¹. The lowest flower production per hectare was obtained from a combination of treatments without applying chicken manure and biochar (control).

References

- [1] ADNAN. 2018. Growth and Yield of cauliflower Pertumbuhan Dan Hasil Kubis Bunga (*Brassica oleraceae*) due to different seedling times and applying various doses of Kompos fertilizer. *J. Research of Agrosamudra*. Vol 5(1): 1-13.

- [2] AMALIA, S., NURDIANA, D., & MAESYAROH, S. S. (2019). Effect of Chicken Manure Dose and Fungus of *Trichoderma* sp. On Growth and Yield of Cauliflower (*Brassica oleracea* var. *Botritis* L.). *Jagros: Journal of Agrotechnology Science*, 3(2), 122–135. <http://dx.doi.org/10.52434/jagros.v3i2.865>
- [3] ALI, M. 2001. Leaching Impact on crop production and health. UPN Press. Surabaya.
- [4] BAHRI. (2006). Effect of sources of Fertilizer on growth and Yield of Cauliflower (*Brassica oleracea* var. *Botritis* L.). Presiding of National Seminar of Horticulture, 307–310. Available on <http://sumbar.litbang.pertanian.go.id/images/pdf/selada.pdf>.
- [5] CHAN, K.Y., VAN ZWIETEN, B.L., MESZAROS, I., DOWNIE, D. AND JOSEPH, S. 2007. Agronomic values of green waste biochars as a soil amendment. *Aust J. of Soil Resource*. 45 (2): 629-634
- [6] CAHYONO, B. 2003. *Cauliflower and Broccoli*. Kanisius Publisher Yogyakarta. Yogyakarta Indonesia
- [7] CORREIA, V.P., RUBZEN M. F. R., BRITO, M., SILVA, O., DAHU, S. G.G., 2022. *Impact of Covid-19 on Vegetable Producers: The Case of Cauliflower and Broccoli Farmers in the Municipality of Aileu*. Research Report. Faculty of Agriculture - National University of Timor Lorosa'e (UNTL) and Alumni Research Support Facilities (arsf ii) - ACIAR
- [8] DELLAH, T. S., 2023. *The Effect of Biochar and Goat Manure on the Growth and Yield of Cauliflower (Brassica oleracea var. Botrytis L.) and the Growth of Weed*. Study Program of Agronomy Department of Agronomy University of Sriwijaya, Indonesia.
- [9] EDDUARD Y. N., CEUNFINB, S., THEODORUS T. M. (2020). Effect of Rice Husk Biochar Dosing and Chicken Residu on Growth and Yield of Cauliflower (*Brassica oleraceae*, L.). *Journal of Conservation Agriculture Dryland*. Center of Dryland Research, Timor University, Kefamenanu, TTU – NTT, Indonesia.
- [10] EVANS, M.R., JACKSON, B.E., POPP, M., SADAHA, S. 2017. Chemical properties of biochar material manufactured from agricultural products common to the southeast United States. *Journal Horticulture Technology*. 27,16–23.
- [11] EVIZAL, R., PRASMATIWI, F. R., 2023. Biochar: Utilization and Practical Application. *Journal of Agrotropica* 22 (1):1. Department of Agrotechnology, Faculty of Agriculture, University of Lampung.
- [12] FITRIANI, M. L. 2009. Cultivation of Cauliflower (*Brassica oleraceae* Var *Botrytis* L.) in Horticulture Seed Center Tawangmangu. Faculty of Agriculture, Sebelas Maret University. Surakarta.
- [13] GANI, A., 2009. Biological Charcoal Potential “Biochar” As a technology component of Dryland productivity rehabilitation. Center of Rice Research, Sukamandi.
- [14] GAO, S., HOFFMAN-KRULL, K., BIDWELL, A.L., DELUCA, T.H. 2016. *Locally Produced Wood*

Biochar Increase Nutrient Retention and Availability in Agricultural Soils of the San Juan Islands, USA. Agriculture, Ecosystems & Environment. 233, 43–54.

- [15] GUNAWAN I., ATAK T., ISNA T. (2021) Optimizing dose of Chicken Manure and NPK on Cauliflower Cultivation (Brassica oleracea var. Botrytis L.) AgriHealth: *Journal of Agri-food, Nutrition and Public Health*. 2(1), 32-38. Sebelas Maret University, Department of Agrotechnology, Faculty of Agriculture Garut University, Garut, Indonesia URL: <https://jurnal.uns.ac.id/agrihealth/article/view/49828>
- [16] HUSNIHUDA, M. I., SARWITRI, R., & SUSILOWATI, Y. E. (2017). Response of growth and Yield of Cauliflower (Brassica oleracea Var. Botrytis, L.) on applying PGPR Bamboo Root and Composition of Media Planting. *Vigor: Journal of Agriculture Science of Tropical and Subtropical*, 2(1), 13–16. Available from <https://jurnal.untidar.ac.id/index.php/vigor/article/view/321>.
- [17] ISDARMANTO. (2009). Effect of Various Organic Fertilizers and Leaf Fertilizer Concentration on the Growth and Yield of Red Chili (*Capsicum annum L.*) in Pot System Cultivation. *Faculty of Agriculture*, Sebelas Maret University, Surakarta.
- [18] LEHMANN, J. AND JOSEPH, S., 2009. *Biochar for environmental management: Science and Technology. Journal of Scientific Research an Academic Publisher*. Earthscan: London 127-143.
- [19] MAP, 2020. *Data statistic of Production of Horticulture*. General Director of Plant Production and Horticulture Timor-Lorosa'e.
- [20] MAYADEWI, N. N. A. (2007). Effect of various Manure and Planting Space on the Growth of Weeds and Yield of Sweet Corn. *Journal Agritrop*, 26(4), 153–159. Available from <https://ojs.unud.ac.id/index.php/agritrop/article/view/3069/2217>
- [21] MUSHIN, 2003. *Applying of Chicken Manure on the Growth and Yield of Cucumber. Research Report Agriculture Faculty Taman Siswa University Padang*.
- [22] PURWONO, 2011. Soil science- Soil fertility. Soil Department Faculty of Agriculture, IPB. Indonesia P 48.
- [23] STEINER, C., W. TEIXEIRA, J. LEHMANN, AND W. ZECH. 2003. Microbial response to charcoal amendments of highly weathered soils and Amazonian Dark Earths in Central Amazonia. Dordrecht: Kluwer Academic Publishers. *J. of Soil Resource*1(1): 196-211
- [24] STEWART, B.A., 1991. *Advance in Soil Science*. Vol. 13. Dryland Agriculture: Agriculture for Sustainability. P 72.
- [25] SUBROTO, 2009. *Fertility and Fertilizing of Agriculture Soil*. Bandung: Pustaka Buana. Bandung

Indonesia. P 28

- [26] SUNARTI, 2014. Inportant Pest and Disease Observation on Lowland Cauliflower (*Brassica oleraceae* var. *Botrytis*). *Journal Agroqua*, 13(2).
- [27] SUN, F., AND S. LU., 2014. Biochars improve aggregate stability, water retention, and pore space properties of clayey soil. *J. Plant Nutrition. Soil Science.*, 177, 26–33.
- [28] SUTANTO, R., 2002. *Agriculture Organic Application, Society and the Development*. Publisher Kanisius Yogyakarta- Indonesia. P 6-98.
- [29] WARNOCK, D. D., J. LEHMANN, T. W. KUYPER, AND M. C. RILLIG. 2007. Mycorrhizal responses to biochar in soil ± concepts and mechanisms. *J. Plant and Soil*. 30 (1): 9-20
- [30] WIDOWATI. L. R., SRI WIDATI, JAENUDIN, U., & HARTATIK, W., 2005. Effect of organic Compost with adding Mineral Materials and Biological fertilizers on Soil Character, Absorb of Mineral nutrients, and Production of Organic Vegetable. *Journal Report of Research Project Agribusiness Development Program, Soil Research Center*.
- [31] WIDOWATI. L. R., 2009. Role of Organic Fertilizers on Fertilizing Efficiency and Nesecity Level of Vegetable Plant in Inseptisols Soil Ciherang. Bogor. *Journal Tropica Soil*, vol 14 p 221-228
- [32] WINANDA A., EFENDI E., SAFRUDDIN F., 2019. Response Applying NPK Grower and Chicken Fese on the Growth and Yield of Onion. *Bernas Agricultural Research Journal*. Department of Agrotechnology Agriculture Faculty, Asahan University. *Journal.una.ac.id*.